Preprint typeset in JHEP style - HYPER VERSION

arXiv:1210.4740v2 [hep-th] 23 Oct 2012

# The Negative Mode for small CDL instantons

---

**I-Sheng Yang**[*]

*IOP and GRAPPA, Universiteit van Amsterdam, Science Park 904, 1090 GL Amsterdam, Netherlands*

Abstract: Under thin-wall approximation, we explicitly construct the $O(4)$ symmetric negative mode for the small Coleman-de Luccia instantons. It turns out to be the same mode that changes the bubble size as for the big instantons. We argue that both mediate vacuum transitions in similar ways and provide the thermal interpretations.

---

[*] isheng.yang@gmail.com

## Contents

1. **Introduction** 1

2. **The Small CDL Instanton** 2

3. **Recovering the Negative Mode** 5

4. **The Thermal Interpretation** 7

5. **Conclusion** 8

## 1. Introduction

Coleman and de Luccia wrote down the instanton solution that became the paradigm of first order phase transitions with gravity [1]. When the critical bubble is much smaller than the de Sitter radius of the parent vacuum, the CDL instanton approaches the Coleman instanton in flat space [2]. In the thin-wall approximation, they have the same negative mode which corresponds to changing the critical bubble size. On the other hand, a continuous parameter change from the above "big" instanton leads to a "small" instanton, as depicted in Fig.1. Such solution is troublesome in two aspects. First, it appears to lose the usual negative mode corresponding to the change of bubble size [3]. Second, having less than half of the parent de Sitter 4-sphere defies the usual interpretation of the phase transition which is nucleating a bubble.

The existence and uniqueness of the negative mode is a criterion for the instantons to mediate vacuum transitions. For the Coleman instanton without gravity, it was proved in [4]. Considering the gauge theory nature of gravity, the CDL instanton needs certain appropriate mode reduction process, otherwise there will be spurious modes [5–7]. A tentative existence plus uniqueness proof was presented in [8][1]. Following the same technique, numerical examples of both big and small instantons are shown to have exactly one negative mode [9, 10]. However, such framework cannot clearly demonstrate which physical deformation the negative mode corresponds to, and it does



**Figure 1:** From left to right we show the flat space instanton solution with a critical bubble, the normal (large) instanton solution between two de Sitter vacua, and the small instanton solution between the same pair of de Sitter vacua. The arrows represent the deformation that changes the bubble size, which corresponds to the negative mode in the first two cases, but a positive mode for the last case.

not distinguish between big and small instantons. Thus the sharp contrast described in the previous paragraph is not fully resolved.

In this paper, we will stick to the thin-wall approximation and explicitly construct the physical deformation corresponding to the negative mode for small CDL instantons. Just like for the big instantons, it still corresponds to the change of bubble size. In light of this, we see no reason to treat them differently. In fact, we will show that in the thermal interpretation [11], it represents the phase transition similarly as how a big instanton represents the reverse transition [12].

The structure of this paper goes like the following. In section 2, we review the basic solution of a thin-wall CDL instanton and the missing negative mode for the small instanton. In section 3 we explicitly demonstrate the existence of this negative mode. In section 4 we discuss how to interpret the small instanton as mediating a phase transition. Finally we summarize and conclude in section 5.

## 2. The Small CDL Instanton

Consider a scalar field with the following potential,

$$V(\phi) = \lambda^2(\phi^2 - \phi_0^2)^2 - \frac{\Delta V \phi}{2\phi_0} \ . \tag{2.1}$$

We have a true vacuum and a false vacuum at

$$\phi_t \approx \phi_0 \ , \qquad \phi_f \approx -\phi_0 \ , \tag{2.2}$$

---

[1] It is tentative because a not-fully-justified analytical continuation for the uniqueness proof, and a possible sign problem of $Q_E$ in the existence proof as pointed out in [9].

with energy difference roughly $\Delta V$, and a domain wall separating them with tension given by

$$\sigma \approx \int_{\phi_f}^{\phi_t} \sqrt{2V} d\phi \ . \tag{2.3}$$

Given that $\Delta V \ll \lambda^2 \phi_0^4$, a false vacuum background can nucleate a thin-wall bubble of true vacuum, which will then expand and complete the phase transition. The rate of this nucleation process is given by (keeping only the exponent)

$$\Gamma_{f\to t} = e^{S_I - S_f} \ , \tag{2.4}$$

$$S = \int L_m \ dx^4 = \int \left( \frac{1}{2} (\partial\phi)^2 + V(\phi) \right) dx^4 \ . \tag{2.5}$$

$S$ stands for the Euclidean action: $S_f$ for the background false vacuum and $S_I$ for the instanton solution that contains a bubble of true vacuum. The 4D Euclidean configuration of the instanton solution is a 3-sphere of domain wall, filled with the true vacuum and surrounded by the false vacuum. One can easily write down the action difference,

$$(S_I - S_f) = 2\pi^2 r^3 \sigma - \frac{\pi^2}{2} r^4 \Delta V \ . \tag{2.6}$$

It has a maximum at

$$r_c = 3\frac{\sigma}{\Delta V} \ . \tag{2.7}$$

Varying $r$ is an unique negative mode that signifies this configuration being the leading saddle point contribution that mediates the phase transition.

Including gravity, the Euclidean action becomes:

$$S = \int \sqrt{g} dx^4 \left( L_m - \frac{M_p^2}{2} \mathcal{R} \right) \ , \tag{2.8}$$

where the matter action is

$$L_m = \frac{1}{2} g_{\mu\nu} \partial^\mu \phi \partial^\nu \phi + V \ . \tag{2.9}$$

For simplicity, in this paper we will focus on the scenario that both vacua has positive energy by adding a constant term to $V$, such that

$$V(\phi_t) = \frac{3M_p^2}{R_t^2} \ , \quad V(\phi_f) = \frac{3M_p^2}{R_f^2} \ . \tag{2.10}$$

The Euclidean action of the false vacuum configuration is simply

$$S_f = \left( V(\phi_f) - \frac{6M_p^2}{R_f^2} \right) \ \mathcal{V}(R_f, \text{full}) \ . \tag{2.11}$$

The two terms in the bracket are the field and gravity contributions, and the last factor stands for the 4-volume of a 4-sphere with radius $R_f$.

The instanton is the matching of two 4-spheres with radii $R_t$ and $R_f$ at some junction radius $r$, where the domain wall resides. Its action is given by

$$S_I = 2\pi^2 r^3 \sigma + L_{g,wall}(R_f, R_t, r) + \left( V(\phi_f) - \frac{6M_p^2}{R_f^2} \right) \mathcal{V}(R_f, r) + \left( V(\phi_t) - \frac{6M_p^2}{R_t^2} \right) \mathcal{V}(R_t, r) \, . \tag{2.12}$$

The second term is the gravitational contribution from the junction[2],

$$L_{g,wall}(R_f, R_t, r) = 6\pi^2 M_p^2 r^2 \left( \pm \sqrt{1 - \frac{r^2}{R_f^2}} \pm \sqrt{1 - \frac{r^2}{R_t^2}} \right) \, . \tag{2.13}$$

It takes the plus sign when the corresponding side is a portion of 4-sphere that contains a full equator 3-sphere, namely the bigger portion, and negative sign for a smaller portion. The same ambiguity appears in $\mathcal{V}(R_f, r)$, which means the volume of a partial 4-sphere bounded by a 3-sphere of radius $r$ and can be either the bigger or the smaller side. Fortunately as shown in [1, 13] that all these ambiguous terms can be combined to show that without ambiguity, the instanton action is extremized at

$$r_e = \frac{r_c}{\sqrt{1 + (R_f^{-2} + R_t^{-2})\frac{r_c^2}{2} + (R_f^{-2} - R_t^{-2})^2 \frac{r_c^4}{16}}} \, , \tag{2.14}$$

where $r_c$ is the critical bubble size in flat space, given by Eq. (2.7).

When $r_c \ll R_f$, the instanton contains a small portion of the true vacuum and a large portion of the false vacuum, $r_e \sim r_c$, and varying $r$ is a negative mode just as in the case without gravity. As we tune the potential to increase $r_c$, for example reducing $\Delta V$, $r_e$ smoothly becomes inversely proportional to $r_c$ instead, at the same time the instanton becomes two smaller-than-half portion of both the true and the false vacuum. These two cases are drawn in Fig.1. The annoying fact is, in the later case, varying $r$ also becomes a positive mode [3].

For this kind of small instanton, its action is still bigger than $S_f$ (and $S_t$), so people tend to believe that it has at least one negative mode[3]. Since the above analysis is restricted to $O(4)$ symmetry and thin-wall approximation, the common intuition is to go beyond either or both of them. Maybe the disappearance of the radial negative mode signifies the emergence of many more subtle negative modes, and condensing them leads to a thick-wall or less symmetric solution that has only one negative mode.

---

[2] Note that this term comes from Eq. (3.6), in which the Racci scalar $\mathcal{R}$ receives a delta function contribution from the second derivative term since the first derivative is discontinuous across the junction. In many literatures, one integrates by part to turn this term into a Gibbons-Hawking (surface) term and a modification to the Einstein-Hilbert (volume) term. We will refrain from doing the same so the physical meaning of each term remains clear.

[3] Of course, it can also be a local minimum isolated from both vacuum configurations. That is typically considered less likely.

However, these suggestions are all motivated by the apparent "disappearance" of the radial negative mode. We will explicitly show that such a mode actually still exists.

## 3. Recovering the Negative Mode

Let us reconsider what happened in the extremizing process in the previous section. At the extremum, we have

$$2\pi^2 r_e^3 \sigma = -\frac{3}{2} L_{g,wall}(R_f, R_t, r_e) \ , \tag{3.1}$$

which is the Israel junction condition [14], namely the integrated Einstein equations across a codimension one delta function. This equality will not hold at any other radius, which means varying the radial mode is in a phase space that does not solve the full equations of motion,

$$\phi'' + 3\frac{\rho'}{\rho}\phi' = \frac{\partial V}{\partial \phi} \ , \tag{3.2}$$

$$\rho'^2 = 1 + \frac{\rho^2}{3M_p^2}\left(\frac{\phi'^2}{2} - V\right) \ , \tag{3.3}$$

$$\rho'' = -\frac{2}{3M_p^2}(\phi'^2 + V) \ , \tag{3.4}$$

with the metric

$$ds^2 = d\xi^2 + \rho^2 d\Omega_3^2 \ . \tag{3.5}$$

However, note that

$$L_{g,wall} = -\frac{M_p^2}{2}\int_{\bar{\xi}-\epsilon}^{\bar{\xi}+\epsilon} \mathcal{R}\rho^3 d\xi \ , \tag{3.6}$$

with $\epsilon \to 0$ and $\rho(\bar{\xi}) = r$. Its value only involves the $\rho''$ term in $\mathcal{R}$. So not solving the junction condition just means not solving Eq. (3.4).

The common lore in semiclassical quantum gravity is that Eq. (3.3), as a first order equation, should be treated as a constraint, similar to gauge fixing in a gauge theory. One should not consider deformations violating this constraint in order to avoid spurious modes [5–7]. On the other hand, Eq. (3.4) is a second order equation, thus a dynamical equation of motion. Just like we should consider modes violating the field equation of motion, Eq. (3.2), it is also fine to violate Eq. (3.4). Both of them will only be satisfied at the critical points.

Without a full theory of quantum gravity, one can always question this semiclassical approach. However that is not the focus of this paper. We will simply do exactly the same thing as Coleman and de Luccia did. **In the variation of CDL radial mode, we go through configurations where the gravity contribution to the junction, Eq. (2.13), does not match the matter contribution from the domain wall tension.** For the same token, we shall be allowed to do the following. **Put a geometric junction where there is no domain wall. It only contributes gravitationally, as if there is a zero tension domain wall.** The constraint equation demands only that in the true (false) vacuum region, the geometry has to be a portion of the 4-sphere with $R_t$ ($R_f$).

With that in mind, let us consider the following solution parameterized by two radii, $r_g$ and $r_w$. At $r_g$ there is a purely geometric junction, and at $r_w$ we have the usual domain wall with tension $\sigma$ separating the true and false vacuum. When the purely geometric junction is in the true vacuum region, we have

$$\begin{aligned} S_I(r_g, r_m) = {} & 2\pi^2 r_m^3 \sigma + L_{g,wall}(R_f, R_t, r_m) + L_{g,wall}(R_t, R_t, r_g) \\ & + \left( V(\phi_f) - \frac{6M_p^2}{R_f^2} \right) \mathcal{V}(R_f, r_m) \\ & + \left( V(\phi_t) - \frac{6M_p^2}{R_t^2} \right) \Big( \mathcal{V}(R_t, r_m \text{ to } r_g) + \mathcal{V}(R_t, r_g) \Big) \, . \end{aligned} \tag{3.7}$$

When it is in the false vacuum region, just switch $t$ and $f$ in the above equation. All the 4-volume functions $\mathcal{V}$ here are referring to the smaller portion without an equator 3-sphere, which should be obvious from Fig.2.

When $r_g = r_m = r_e$, this is exactly the small CDL instanton as described in the previous section. Now we can fix $r_g = r_e$ and starts to vary $r_m$ to either side, as shown in Fig.2. We see that the action always decreases[4]. We can further shrink $r_m$ to zero to eliminate the true or false vacuum portion, and smooth out the purely geometric junction. The action is strictly decreasing during the entire process and recovers $S_t$ or $S_f$.

The analysis of radial mode in [3] was restricted to a single geometric junction that always sticks with the domain wall. This unnecessary restriction led to a bias that for

---

[4] We skipped the equations or numerical plots here since one can easily observe the following: the change of action is dominated by the domain wall contribution, which is shrinking in both directions of this deformation.



**Figure 2:** The top middle figure is a small CDL instanton. We can deform the domain wall (dashed line) away from its critical position to either left or right while leaving a purely geometric junction (dotted line) behind. This is a negative mode as the action decreases in both directions. We can further shrink the domain wall and smooth out the junction to recover the true or false vacuum 4-sphere—solutions without negative modes.

a small instanton, changing the bubble size also changes the total size of the entire instanton. What we did was to remove that bias, such that changing the bubble size really means increasing/decreasing the region of true/false vacuum. Then expectedly, the negative mode reappears.

## 4. The Thermal Interpretation

Brown and Weinberg [11] provided a very accurate picture to interpret how a large CDL instanton mediates the vacuum transition. Instead of taking the Euclidean 4-sphere as the global geometry, they described it as a horizon 3-volume times a compact coordinate from the finite temperature. One side of the equator is the horizon volume before tunneling, and the other side is the same horizon volume after tunneling.

This interpretation clarified a few confusions. For example, without an exactly thin wall, the "false vacuum region" of the instanton will not be identical to the same portion of the false vacuum 4-sphere. If one takes the instanton as a global geometry, it is unsatisfying that nucleating a bubble requires changes far away, out of causal contact from the bubble. In the thermal interpretation this has a clear explanation. With nonzero temperature, the transition is not purely quantum, but always thermally assisted, as depicted in Fig.3. The horizon volume of the false vacuum always needs to be thermally excited, even just a little bit, to the configuration that is the left hand side slicing of an instanton, then the quantum tunneling starts.

Note that not only the field configuration of the instanton is slightly away from the pure vacuum, so is the geometry. This is also straight forward since the gravitational back reaction from a non-vacuum state leads to a non-vacuum geometry[5]. Realizing this fact also means that we can accept the reverse tunneling being mediated by the same instanton, just in the reverse direction. It is a dramatic fluctuation from the true vacuum to the initial condition of this reverse tunneling, in terms of both the field configuration and the geometry. From the horizon volume of the true vacuum, the fluctuation leads to a bubble of true vacuum surrounded by false vacuum regions, and a much reduced horizon size[6]. But that is just what is has to be and most of the suppression in the tunneling rate is indeed a thermal factor.

With these in mind, the small instanton mediates tunneling in the same way, only that both directions require a dramatic thermal fluctuation. From a horizon volume of a vacuum, we need a thermal fluctuation up to a smaller volume surrounded by a domain wall, as shown in Fig.4.

## 5. Conclusion

We have shown that under thin-wall approximation and the same principle of semi-classical quantum gravity, the small CDL intanton has the same radial negative mode as the large CDL instanton. This was not known before due to an over-restriction of the geometry that we removed by allowing a purely geometric junction. Although a full analysis including thick-wall effects and less symmetries is still lacking, our result agrees with the numerical thick wall example in [9, 10] and we believe the difference between the big and small instantons is eliminated. The big instantons, being similar to the Coleman instanton in flat space, has been widely accepted as the correct saddle point for the tunneling. The same should be true for the small instantons. We provided the thermal interpretation for upward and downward tunnelings mediated by big and small instantons.

## Acknowledgments

I am especially grateful that Erick Weinberg has planted this problem deeply in my

---

[5] Therefore, it remains to be clarify as how the equator of a instanton is cut into the initial and final "horizon volume". We will take the mathematically obvious choice: the boundary is the maximum 2-sphere. The physical reason behind this choice might worth further investigations.

[6] We thank Adam Brown for a brief discussion about this.



**Figure 3:** The downward (red, left to right) and upward (blue, right to left) vacuum transitions mediated by the large CDL instanton. Both processes are shown by two dashed-arch-arrow, one for the thermal fluctuation from the initial horizon volume to the configuration of the corresponding (almost) semi-3-sphere on the instanton geometry, the other for the tunneling to the other side of the instanton geometry. On the instanton geometry, we use the maximum 2-sphere to separate the two horizon volumes before and after the tunneling. The downward transition obviously involves a smaller thermal fluctuation.



**Figure 4:** The same figure for a small CDL instanton. Here, both downward and upward tunnelings involve some dramatic thermal fluctuations to begin with.

mind. I also thank Adam Brown, Bartek Czech, Ben Freivogel and Jean-Luc Lehners for stimulating discussions. This work is part of the research program of the Foundation for Fundamental Research on Matter (FOM), which is part of the Netherlands Organization for Scientific Research (NWO).

## References

[1] S. Coleman and F. D. Luccia, "Gravitational effects on and of vacuum decay," *Phys. Rev. D* **21** (1980) 3305–3315.

[2] S. Coleman, "THE FATE OF THE FALSE VACUUM. 1. SEMICLASSICAL THEORY," *Phys. Rev. D* **15** (1977) 2929–2936.

[3] K. Marvel and N. Turok, "Horizons and Tunneling in the Euclidean False Vacuum," `arXiv:0712.2719 [hep-th]`.

[4] S. R. Coleman, "QUANTUM TUNNELING AND NEGATIVE EIGENVALUES," *Nucl.Phys.* **B298** (1988) 178.

[5] T. Tanaka and M. Sasaki, "False vacuum decay with gravity: Negative mode problem," *Prog.Theor.Phys.* **88** (1992) 503–528.

[6] T. Tanaka, "The No - negative mode theorem in false vacuum decay with gravity," *Nucl.Phys.* **B556** (1999) 373–396, `gr-qc/9901082`.

[7] G. V. Lavrelashvili, "Negative mode problem in false vacuum decay with gravity," *Nucl.Phys.Proc.Suppl.* **88** (2000) 75–82, `gr-qc/0004025`.

[8] A. Khvedelidze, G. V. Lavrelashvili, and T. Tanaka, "On cosmological perturbations in closed FRW model with scalar field and false vacuum decay," *Phys.Rev.* **D62** (2000) 083501, `gr-qc/0001041`.

[9] L. Battarra, G. Lavrelashvili, and J.-L. Lehners, "Negative Modes of Oscillating Instantons," `1208.2182`.

[10] G. Lavrelashvili, "The Number of negative modes of the oscillating bounces," *Phys.Rev.* **D73** (2006) 083513, `gr-qc/0602039`.

[11] A. R. Brown and E. J. Weinberg, "Thermal derivation of the Coleman-De Luccia tunneling prescription," *Phys. Rev.* **D76** (2007) 064003, `arXiv:0706.1573 [hep-th]`.

[12] K. Lee and E. J. Weinberg, "Decay of the true vacuum in curved space-time," *Phys. Rev. D* **36** (Aug, 1987) 1088–1094.

[13] S. Parke, "Gravity and the decay of the false vacuum," *Physics Letters B* **121** (1983) no. 5, 313 – 315.

[14] W. Israel, "Singular hypersurfaces and thin shells in general relativity," *Nuovo Cim.* **B44S10** (1966) 1.